



LAVANDULA angustifolia 'Hidcote Superior'


True Lavender, English Lavender

Item No.: LA076

Also available as: GOLD NUGGET SEED ®

Portion Price (sufficient for 50-100 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
3,00€	6,00€	48,00€	384,00€	-	-

Plant Description

Life Cycle	Perennial
Family	Lamiaceae
Origin	Garden origin
Special Features	Our own production of 'Hidcote Superior' ensures a predictable and constant quality and is available exclusively from Jelitto. Even and constant quality. Great for containers, in the border or for edging.
Historical	Introduced by JELITTO PERENNIAL SEED in 2001.
Basic Colour	(violet)
Flower Colour	deep violet blue
Natural Flowering Period	July - September
Winter Hardiness Zones	Z5 - Z9
Foliage	whitish-gray, lanceolate (or narrower)
Growth Habit	compact / shrubby
Height with Flowers	40 cm
Spacing between Plants	50 cm
Soil Requirements	dry / well-drained
Location	
Characteristics	ornamental foliage plant
Usage	honey-bee food plant / for the rock garden

Cultivation

Grams per 1000 seeds	0.71429 Gram
Seeds per Gram	1400 (does not correspond to the number of plants!)



Gram to get 1000 plants	5 Gram (if sown directly into pots etc. you will need a larger quantity)
Sowing Rates/Trays	4 - 8 per cell
Plug tray recommended size(s)	72 / 128
Sowing Direction	(9) These seeds germinate rapidly depending on species and origin. If germination does not occur after 3–4 weeks a cooling period of 2–4 weeks is recommended.

Scheduling

Best Sowing Date	late autumn - early spring (northern Hemisphere, Field condition)
Germination to Transplant	4 - 8 weeks
Transplanting to Potting	6 - 10 weeks
Cutting back at Transplanting	Cut-back once to keep more compact.

Growing On

Container Size(s)	1 plug per 8/9 cm (3 1/2") / 1-2 plugs per 11/12 cm (4 1/2")
Vernalization	There is no current research on vernalization but a prudent recommendation for any perennial would be 6-12 weeks (a few might need 15 weeks!) at an average daily temperature of 40°F (5°C). Exposure to cold may not be necessary for flowering but might improve quality.
Forcing	An obvious place to experiment - following vernalization - would be raising daytime temperatures to 60° - 65°F (15° - 17°C). Provide 16 hours of continuous lighting. During the short days of winter, provide a night interruption lighting of 4 hours between 10:00 p.m. and 2:00 a.m.
Fertilizer	Light (100-150 ppm)