



## CAMPANULA persicifolia 'Grandiflora Alba'



[Weisse Glocke, White Bells]

Willow Bell, Peach Bells, Peach Leaf

Item No.: CA182

Portion Price (sufficient for 50-100 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
2,00€	4,00€	32,00€	256,00€	-	-

### Plant Description

<b>Life Cycle</b>	Perennial
<b>Family</b>	Campanulaceae
<b>Origin</b>	Europe, North Africa
<b>Historical</b>	The species has been cultivated since the 16th century.
<b>Basic Colour</b>	(white / cream)
<b>Flower Colour</b>	white
<b>Natural Flowering Period</b>	June - July
<b>Winter Hardiness Zones</b>	Z3 - Z7
<b>Foliage</b>	leathery, dark, strap-shaped basal leaves
<b>Growth Habit</b>	erect
<b>Height with Flowers</b>	100 cm
<b>Spacing between Plants</b>	45 cm
<b>Soil Requirements</b>	dry / well-drained
<b>Location</b>	 
<b>Usage</b>	honey-bee food plant / suitable for cutting

### Cultivation

<b>Grams per 1000 seeds</b>	0.06061 Gram
<b>Seeds per Gram</b>	16500 (does not correspond to the number of plants!)
<b>Gram to get 1000 plants</b>	0.5 Gram (if sown directly into pots etc. you will need a larger quantity)
<b>Sowing Rates/Trays</b>	8 per cell
<b>Plug tray recommended size(s)</b>	128 / 288



**Sowing Direction**

(16) See 15 with the exception of germination being slower and more irregular. This poses no problems.

(15) Rapidly germinating, keep seed in constant moisture (not wet) with temperatures of about +20°C [68°F]. Seeds must be covered thinly. Do not cover very small seeds, but tightly press into the earth. Keep in cooler conditions after germination occurs.

**Scheduling**

**Best Sowing Date**

Anytime (northern Hemisphere, Field condition)

**Sowing to Germination**

3 - 4 weeks

**Germination to Transplant**

4 - 6 weeks

**Transplanting to Potting**

6 - 8 weeks

**Cutting back at Transplanting**

Not Necessary.

**Growing On**

**Container Size(s)**

1-2 plugs per 11/12 cm (4 1/2") / 2-3 plugs per 15 cm (6")

**Vernalization**

A prudent recommendation would be to provide 6-12 weeks at an average daily temperature of 40°F (5°C). Exposure to cold may not be necessary for flowering but might improve quality.

**Forcing**

Provide 16 hours of continuous lighting. During the short days of winter, provide a night interruption lighting of 4 hours between 10:00 p.m. and 2:00 a.m. 6-8 weeks to flower.

**Fertilizer**

Medium (150-200 ppm)