



NECTAROSCORDUM siculum



(ALLIUM)

Mediterranean Bells

Item No.: NA015

Portion Price (sufficient for 50-100 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
3,00€	6,00€	48,00€	384,00€	-	-

Plant Description

Life Cycle	Perennial
Family	Alliaceae
Special Features	Loose umbels.
Basic Colour	(green)
Flower Colour	greenish-white with pink touch
Natural Flowering Period	May - June
Winter Hardiness Zones	Z5 - Z8
Growth Habit	bulbous
Height with Flowers	80 cm
Soil Requirements	average
Location	 
Usage	suitable for cutting

Cultivation

Grams per 1000 seeds	5.71429 Gram
Seeds per Gram	175 (does not correspond to the number of plants!)
Gram to get 1000 plants	15 Gram (if sown directly into pots etc. you will need a larger quantity)

Sowing Direction

(1) Cold-germinators are still referred to as frost-germinators, although this isn't quite correct. The sowing must be kept warm (about +18 to +22°C) [about 64 to 72°F] and moist for the first 2–4 weeks. After this period the sowing must be kept at a cold temperature (between –4 and +4°C) [between 25 and 39°F] for another 4–6 weeks. Colder temperatures of –5°C [23°F] are only advantageous for most species of the Ranunculus family. It is not so important if the temperature is higher or lower during the cooling period, but the cooling period has to be prolonged because the synthesis of the germination inducer, hormon-like acid, slows down or comes to a standstill. It is beneficial to cover the sowing with snow during the cooling-period. The temperature below it usually keeps in the optimum range of –4 to 0°C [25 to 32°F]. The sowing is kept moist, and the melting snow helps to destroy the shell, which is advantageous for the germinating seedling. After this cooling-period the sowing may not be immediately exposed to high temperatures. The most effective temperatures are between +5 to +12°C [41 to 54°F], even if germination has started. The best location for this sowing, even in March, April and May, is the open field, the cold frame or a cold greenhouse.