



## BELAMCANDA flabellata 'Hello Yellow'


(flava)

Dwarf Blackberry Lily

Item No.: BA027

Portion Price (usually sufficient for approx. 50 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
5,80€	5,80€	46,00€	368,00€	-	-

### Plant Description

<b>Life Cycle</b>	Perennial
<b>Family</b>	Iridaceae
<b>Origin</b>	Asia
<b>Special Features</b>	Charming large, yellow blooms and decorative seed pods with shiny blackberry-like seeds. Beautiful cut flower - fresh or dried. Drought tolerant/ for xeriscaping/ waterwise design. Deer resistant. Moderate self-sowing at suitable locations. Easy to grow.
<b>Basic Colour</b>	(yellow / gold)
<b>Flower Colour</b>	golden
<b>Natural Flowering Period</b>	July - September
<b>Winter Hardiness Zones</b>	Z5 - Z9
<b>Foliage</b>	entire, sword-like, green
<b>Growth Habit</b>	erect / rhizomatous / compact
<b>Height with Flowers</b>	40 cm
<b>Spacing between Plants</b>	30 cm
<b>Soil Requirements</b>	well-drained / average
<b>Location</b>	
<b>Characteristics</b>	decorative fruit
<b>Usage</b>	suitable for cutting

### Cultivation

<b>Grams per 1000 seeds</b>	25 Gram
<b>Seeds per Gram</b>	40 (does not correspond to the number of plants!)



<b>Gram to get 1000 plants</b>	40 Gram (if sown directly into pots etc. you will need a larger quantity)
<b>Sowing Rates/Trays</b>	2 - 3 per cell
<b>Plug tray recommended size(s)</b>	72 / 128
<b>Sowing Direction</b>	<p>(1) Cold-germinators are still referred to as frost-germinators, although this isn't quite correct. The sowing must be kept warm (about +18 to +22°C) [about 64 to 72°F] and moist for the first 2–4 weeks. After this period the sowing must be kept at a cold temperature (between –4 and +4°C) [between 25 and 39°F] for another 4–6 weeks. Colder temperatures of –5°C [23°F] are only advantageous for most species of the Ranunculus family. It is not so important if the temperature is higher or lower during the cooling period, but the cooling period has to be prolonged because the synthesis of the germination inducer, hormon-like acid, slows down or comes to a standstill. It is beneficial to cover the sowing with snow during the cooling-period. The temperature below it usually keeps in the optimum range of –4 to 0°C [25 to 32°F]. The sowing is kept moist, and the melting snow helps to destroy the shell, which is advantageous for the germinating seedling. After this cooling-period the sowing may not be immediately exposed to high temperatures. The most effective temperatures are between +5 to +12°C [41 to 54°F], even if germination has started. The best location for this sowing, even in March, April and May, is the open field, the cold frame or a cold greenhouse.</p> <p>(7) These seeds must be covered with a layer seed kernel diameter. Keep daytime temperatures at approximately +20°C [68°F], and keep the moisture constant, for Cyclamen, however, +18°C [64°F] is recommended. The seeds must be kept in the light and moderately warm after germination.</p>

## Scheduling

<b>Best Sowing Date</b>	late autumn - early spring (northern Hemisphere, Field condition)
<b>Sowing to Germination</b>	8 - 10 weeks
<b>Germination to Transplant</b>	4 - 6 weeks
<b>Transplanting to Potting</b>	6 - 8 weeks

## Growing On

<b>Container Size(s)</b>	1-2 plugs per 11/12 cm (4 1/2") / 2-3 plugs per 15 cm (6")
<b>Fertilizer</b>	Medium (150-200 ppm)