



CORYDALIS acaulis


(PSEUDOFUMARIA alba ssp. acaulis)

Creamy white Fumewort

Item No.: CA652

Portion Price (sufficient for 50-100 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
4,50€	18,00€	-	-	-	-

Plant Description

Life Cycle	Perennial
Family	Papaveraceae
Origin	Southern and Central Europe: stone walls, cracks.
Special Features	Creamy white Fumewort has a firm growth, strong leaves and a good stability. The racemose inflorescences are embedded in the bluish-green, fern-like foliage until the onset of winter. Excellent for dry stone walls and cracks. Aromatic scent. Attractive for butterflies and other pollinators. Drought tolerant/ for xeriscaping/ water-wise design. Suitable for natural landscaping. Moderate self-sowing at suitable locations.
Basic Colour	(white / cream)
Flower Colour	creamy white
Natural Flowering Period	May - October
Flowering	First year flowering
Winter Hardiness Zones	Z5 - Z8
Foliage	bluish green, feathery, fern like, 3-parted, lobed, wintergreen
Growth Habit	compact / wintergreen
Height with Flowers	25 cm
Spacing between Plants	15 cm
Soil Requirements	dry / well-drained / average
Location	
Usage	for the rock garden

Cultivation



Grams per 1000 seeds	1.25 Gram
Seeds per Gram	800 (does not correspond to the number of plants!)
Gram to get 1000 plants	3 Gram (if sown directly into pots etc. you will need a larger quantity)
Sowing Direction	(19) To stop the germination inhibition give seeds a warm compost (about +22°C) [about 72°F] with constant humidity for at least 6 weeks. Then keep cold (-4 to +4°C) [25 to 39°F] for 6–8 weeks. Usually, the germination starts at +4°C [39°F]. Raise temperatures gradually up to +10°C [50°F] until germination is completed. If the warm or cold period was not long enough the seeds will not germinate until the following year. They will then need another warm period (summer) and a cold period (winter). Well-known examples: Cimicifuga and Helleborus.

Scheduling

Best Sowing Date	Anytime (northern Hemisphere, Field condition)
Cutting back at Transplanting	Not Necessary.

Growing On

Container Size(s)	1 plug per 8/9 cm (3 1/2")
Fertilizer	Light (100-150 ppm)