




TANACETUM macrophyllum

Wolfie's Choice

Item No.: CA434

Portion Price (usually sufficient for approx. 50 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
2,00€	4,20€	36,00€	-	-	-

Plant Description

Life Cycle	Perennial
Family	Asteraceae
Origin	Central Europe and Southern Russia
Special Features	Bold, upright appearance, topped with broad 15 cm, flat-topped white blossoms. Modest.
Historical	A favorite of legendary landscape architect, Wolfgang Oehme of Washington, D.C.
Basic Colour	(white / cream)
Flower Colour	white
Natural Flowering Period	June - July
Winter Hardiness Zones	Z5 - Z7
Foliage	fern-like leaves which look more like those of an ACHILLEA
Growth Habit	erect
Height with Flowers	120 cm
Spacing between Plants	120 cm
Soil Requirements	dry / well-drained / average
Location	
Usage	suitable for cutting

Cultivation

Grams per 1000 seeds	0.24096 Gram
Seeds per Gram	4150 (does not correspond to the number of plants!)
Gram to get 1000 plants	3 Gram (if sown directly into pots etc. you will need a larger quantity)



Sowing Rates/Trays	8 per cell
Plug tray recommended size(s)	128 / 288
Sowing Direction	(16) See 15 with the exception of germination being slower and more irregular. This poses no problems. (15) Rapidly germinating, keep seed in constant moisture (not wet) with temperatures of about +20°C [68°F]. Seeds must be covered thinly. Do not cover very small seeds, but tightly press into the earth. Keep in cooler conditions after germination occurs.

Scheduling

Best Sowing Date	Anytime (northern Hemisphere, Field condition)
Sowing to Germination	2 - 3 weeks
Germination to Transplant	4 - 8 weeks
Transplanting to Potting	6 - 10 weeks
Cutting back at Transplanting	Cut back 1-2 times; once at transplanting; once again 4 weeks later to keep more compact.

Growing On

Container Size(s)	1-2 plugs per 11/12 cm (4 1/2")
Vernalization	A prudent recommendation would be to provide 6-12 weeks at an average daily temperature of 40°F (5°C). Exposure to cold may not be necessary for flowering but might improve quality.
Forcing	Provide 16 hours of continuous lighting. During the short days of winter, provide a night interruption lighting of 4 hours between 10:00 p.m. and 2:00 a.m. 6-8 weeks to flower.
Fertilizer	Medium (150-200 ppm)