



TRICYRTIS Hirta-Hybr. 'Miyazaki'


Toad Lily

Item No.: TA192

Also available as: GOLD NUGGET SEED ®

Portion Price (sufficient for 50-100 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
6,80€	42,00€	340,00€	2.700,00€	-	-

Plant Description

Life Cycle	Perennial
Family	Convallariaceae
Origin	Japan
Special Features	Orchid-like autumn blooms. Lovely cut flower for autumn. Flowers emerge from the leaf axils and the stem ends. Elegant arching, sturdy stems.
Basic Colour	(white / cream)
Flower Colour	white with intriguing lilac spots
Natural Flowering Period	September - November
Winter Hardiness Zones	Z4 - Z8
Foliage	medium green, hairy, leaves clasp the stems
Growth Habit	arching / sturdy
Height with Flowers	80 cm
Spacing between Plants	60 cm
Soil Requirements	moist
Location	
Usage	suitable for cutting

Cultivation

Grams per 1000 seeds	0.2439 Gram
Seeds per Gram	4100 (does not correspond to the number of plants!)
Gram to get 1000 plants	1 Gram (if sown directly into pots etc. you will need a larger quantity)
Sowing Rates/Trays	4 per cell
Plug tray recommended size(s)	128 / 288



Sowing Direction

(9) These seeds germinate rapidly depending on species and origin. If germination does not occur after 3–4 weeks a cooling period of 2–4 weeks is recommended.

Scheduling

Best Sowing Date

late autumn - early spring (northern Hemisphere, Field condition)

Germination to Transplant

4 - 6 weeks

Transplanting to Potting

6 - 8 weeks

Cutting back at Transplanting

Not Necessary.

Growing On

Container Size(s)

1-2 plugs per 11/12 cm (4 1/2") / 2-3 plugs per 15 cm (6")

Vernalization

A prudent recommendation would be to provide 6-12 weeks at an average daily temperature of 40°F (5°C). Exposure to cold may not be necessary for flowering but might improve quality.

Forcing

There has been no research, but an obvious place to experiment - following vernalization - would be raising daytime temperatures to 60° - 65°F (15° - 17°C). Provide 16 hours of continuous lighting. During the short days of winter, provide a night interruption lighting of 4 hours between 10:00 p.m. and 2:00 a.m. Some later flowering species can be forced in 14 - 16 weeks and perhaps sooner at warmer temperatures. Further experiments are warranted.

Fertilizer

Medium (150-200 ppm)