



## TRICYRTIS Hirta-Hybr. 'Miyazaki'


Toad Lily

Item No.: TA192

Also available as: GOLD NUGGET SEED ®

Portion Price (sufficient for 50-100 plants)	1g Price (0.1-9.9g)	10g Price (10-99.9g)	100g Price (100-999.9g)	1000g Price (1000-9999.9g)	10000g Price (10000-99999.9g)
6,80€	42,00€	340,00€	2.700,00€	-	-

### Plant Description

<b>Life Cycle</b>	Perennial
<b>Family</b>	Convallariaceae
<b>Origin</b>	Japan
<b>Special Features</b>	Orchid-like autumn blooms. Lovely cut flower for autumn. Flowers emerge from the leaf axils and the stem ends. Elegant arching, sturdy stems.
<b>Basic Colour</b>	(white / cream)
<b>Flower Colour</b>	white with intriguing lilac spots
<b>Natural Flowering Period</b>	September - November
<b>Winter Hardiness Zones</b>	Z4 - Z8
<b>Foliage</b>	medium green, hairy, leaves clasp the stems
<b>Growth Habit</b>	stoloniferous / sturdy
<b>Height with Flowers</b>	80 cm
<b>Spacing between Plants</b>	60 cm
<b>Soil Requirements</b>	moist / average
<b>Location</b>	
<b>Usage</b>	suitable for cutting

### Cultivation

<b>Grams per 1000 seeds</b>	0.24691 Gram
<b>Seeds per Gram</b>	4050 (does not correspond to the number of plants!)
<b>Gram to get 1000 plants</b>	1 Gram (if sown directly into pots etc. you will need a larger quantity)
<b>Sowing Rates/Trays</b>	4 per cell
<b>Plug tray recommended size(s)</b>	128 / 288



**Sowing Direction**

(9) These seeds germinate rapidly depending on species and origin. If germination does not occur after 3–4 weeks a cooling period of 2–4 weeks is recommended.

**Scheduling**

**Best Sowing Date**

late autumn - early spring (northern Hemisphere, Field condition)

**Germination to Transplant**

4 - 6 weeks

**Transplanting to Potting**

6 - 8 weeks

**Cutting back at Transplanting**

Not Necessary.

**Growing On**

**Container Size(s)**

1-2 plugs per 11/12 cm (4 1/2") / 2-3 plugs per 15 cm (6")

**Vernalization**

A prudent recommendation would be to provide 6-12 weeks at an average daily temperature of 40°F (5°C). Exposure to cold may not be necessary for flowering but might improve quality.

**Forcing**

There has been no research, but an obvious place to experiment - following vernalization - would be raising daytime temperatures to 60° - 65°F (15° - 17°C). Provide 16 hours of continuous lighting. During the short days of winter, provide a night interruption lighting of 4 hours between 10:00 p.m. and 2:00 a.m. Some later flowering species can be forced in 14 - 16 weeks and perhaps sooner at warmer temperatures. Further experiments are warranted.

**Fertilizer**

Medium (150-200 ppm)